

CARDINALITIES OF k -DISTANCE SETS IN MINKOWSKI SPACES

K. J. SWANEPOEL

ABSTRACT. A subset of a metric space is a k -distance set if there are exactly k non-zero distances occurring between points. We conjecture that a k -distance set in a d -dimensional Banach space (or *Minkowski space*), contains at most $(k+1)^d$ points, with equality iff the unit ball is a parallelotope. We solve this conjecture in the affirmative for all 2-dimensional spaces and for spaces where the unit ball is a parallelotope. For general spaces we find various weaker upper bounds for k -distance sets.

1. INTRODUCTION

A subset S of a metric space is a k -distance set if there are exactly k non-zero distances occurring between points of S . We also call a 1-distance set an *equilateral set*. In this paper we find upper bounds for the cardinalities of k -distance sets in *Minkowski spaces*, i.e. finite-dimensional Banach spaces (see Theorems 1 to 5), and make a conjecture concerning tight upper bounds.

In Euclidean spaces k -distance sets have been studied extensively; see e.g. [13, 14, 19, 15, 1, 2, 4, 5, 3, 9, 11, 10, 18, 24], and the books [22] and [12, sections F1 and F3].

For general d -dimensional Minkowski spaces it is known that the maximum cardinality of an equilateral set is 2^d , with equality iff the unit ball of the space is a parallelotope, and that if $d \geq 3$, there always exists an equilateral set of at least 4 points [23]. It is unknown whether there always exists an equilateral set of $d+1$ points; see [20, 21] and [25, p. 129, p. 308 problem 4.1.1]. However, Brass [7] recently proved that for each n there is a $d = d(n)$ such that any d -dimensional Minkowski space has an equilateral set of at least n points. See [17] for problems on equilateral sets in ℓ_p spaces.

Equilateral sets in Minkowski spaces have been used in [20] to construct energy-minimizing cones over wire-frames. See also [21].

As far as we know, k -distance sets for $k \geq 2$ have not been studied in spaces other than euclidean.

Our main results are the following.

Theorem 1. *If the unit ball of a d -dimensional Minkowski space is a parallelotope, then a k -distance set in X has cardinality at most $(k+1)^d$. This bound is tight.*

Theorem 2. *Given any set S of n points in a d -dimensional Minkowski space with a parallelotope as unit ball, there exists a point in S from which there are at least $\lceil n^{1/d} \rceil - 1$ distinct non-zero distances to points in S . This bound is tight.*

Theorem 3. *The cardinality of a k -distance set in a 2-dimensional Minkowski space is at most $(k+1)^2$, with equality iff the space has a parallelogram as unit ball.*

Date: 1997.

Key words and phrases. Minkowski space. Erdős distance problem. Equilateral set. k -distance set. Few-distance set.

Theorem 4. *Given any set of n points in a 2-dimensional Minkowski space, there exists a point in S from which there are at least $\lceil n^{1/2} \rceil - 1$ distinct non-zero distances to points in S .*

Theorem 5. *The cardinality of a k -distance set in a d -dimensional Minkowski space is at most $\min(2^{kd}, (k+1)^{(11^d-9^d)/2})$.*

In the light of Theorems 1 and 3 and the results of [23], we make the following

Conjecture 1. *The cardinality of a k -distance set in any d -dimensional Minkowski space is at most $(k+1)^d$, with equality iff the unit ball is a parallelotope.*

As mentioned above, [23] shows that this conjecture is true for $k = 1$. By Theorem 3 the conjecture is true if $d = 2$, and by Theorem 1 if the unit ball is a parallelotope.

In the sequel, $(\mathbb{R}^d, \|\cdot\|)$ is a d -dimensional Minkowski space with norm $\|\cdot\|$, $B(x, r)$ is the closed ball with centre x and radius $r > 0$, and $B := B(0, 1)$ the *unit ball* of the space. Recall that two d -dimensional Minkowski spaces are isometric iff their unit balls are affinely equivalent (by the Mazur-Ulam Theorem; see e.g. [25, Theorem 3.1.2]). In particular, a Minkowski space has a parallelotope as unit ball iff it is isometric to $(\mathbb{R}^d, \|\cdot\|_\infty)$, where $\|(\lambda_1, \lambda_2, \dots, \lambda_d)\|_\infty := \max_{i=1, \dots, d} |\lambda_i|$.

We define a *cone* (or more precisely, an *acute cone*) P to be a convex set in $\mathbb{R}^d$ that is positively homogeneous (i.e., for any $x \in P$ and $\lambda \geq 0$ we have $\lambda x \in P$) and satisfies $P \cap (-P) = \{0\}$. Recall that such a cone defines a partial order on $\mathbb{R}^d$ by $x \leq y \iff y - x \in P$.

We denote the cardinality of a set S by $\#S$.

For measurable $S \subseteq \mathbb{R}^d$, let $\text{vol}(S)$ denote the Lebesgue measure of S . For later reference we state Lyusternik's version of the Brunn-Minkowski inequality (see [8, Theorem 8.1.1]).

Lemma 1. *If $A, B \subseteq \mathbb{R}^d$ are compact, then*

$$\text{vol}(A + B)^{1/d} \geq \text{vol}(A)^{1/d} + \text{vol}(B)^{1/d}.$$

If equality holds and $\text{vol}(A), \text{vol}(B) > 0$, then A and B are convex bodies such that $A = v + \lambda B$ for some $\lambda > 0$ and $v \in \mathbb{R}^d$. $\square$

2. PROOFS

Proof of Theorem 1. We may assume without loss of generality that the space is $(\mathbb{R}^d, \|\cdot\|_\infty)$. We introduce partial orders on $\mathbb{R}^d$ following Blokhuis and Wilbrink [6]. For each $i = 1, \dots, d$, let $\leq_i$ be the partial order with cone

$$P_i = \{(\lambda_1, \dots, \lambda_d) \in \mathbb{R}^d : \max_{j=1, \dots, d} |\lambda_j| = \lambda_i\}.$$

For each x in a k -distance set S , let $h_i(x)$ be the length of the longest descending $\leq_i$ -chain starting with x , i.e. $h_i(x)$ is the largest h such that there exist $x_1, x_2, \dots, x_h \in S$ for which $x >_i x_1 >_i x_2 >_i \dots >_i x_h$.

Since $\bigcup_{i=1}^d (P_i \cup -P_i) = \mathbb{R}^d$, for all distinct $x, y \in \ell_\infty^d$ there exists i such that $x <_i y$ or $y <_i x$. Exactly as in [6], it follows that the mapping $x \mapsto (h_1(x), \dots, h_d(x))$ is injective, and thus $\#S \leq (h+1)^d$, where

$$h := \max_{x \in S, i=1, \dots, d} h_i(x).$$

It remains to show that $h \leq k$. Suppose not. Then for some $x \in S$ and some i there exist $x_1, \dots, x_{k+1} \in S$ such that $x >_i x_1 >_i \dots >_i x_{k+1}$. Since S is a k -distance set, $\|x - x_m\|_\infty = \|x - x_n\|_\infty$ for some $1 \leq m < n \leq k+1$. Also, $x - x_m, x - x_n \in P_i$. Now note that if $\|a\|_\infty = \|b\|_\infty$ with $a, b \in P_i, a \neq b$, then a

and b are $\leq_i$ -incomparable; in particular, $b - a \notin P_i$. Therefore, $x_m - x_n \notin P_i$, a contradiction.

The set $\{0, 1, \dots, k\}^d$ is a k -distance set of cardinality $(k+1)^d$. Note that it is not difficult to see that in fact the only k -distance sets of cardinality $(k+1)^d$ are of the form $S = a + \lambda\{0, 1, \dots, k\}^d$ for some $a \in \mathbb{R}^d$ and $\lambda > 0$. $\square$ $\square$

Proof of Theorem 2. Consider the mapping $x \mapsto (h_1(x), \dots, h_d(x))$ in the proof of Theorem 1. If h is the length of the longest $\leq_i$ -chain over all i , then $n \leq (h+1)^d$. Thus there is a $\leq_i$ -chain $x_0 >_i x_1 >_i \dots >_i x_h$ of length $h \geq \lceil n^{1/d} \rceil - 1$. By the last paragraph of the proof of Theorem 1, the distances $\rho(x_0, x_j), j = 1, \dots, h$ are all distinct.

Any $S \subseteq \mathbb{R}^d$ such that

$$\{0, 1, \dots, \lceil n^{1/d} \rceil - 2\}^d \subsetneq S \subseteq \{0, 1, \dots, \lceil n^{1/d} \rceil - 1\}^d,$$

has exactly $\lceil n^{1/d} \rceil - 1$ distinct distances in the norm $\|\cdot\|_\infty$. $\square$ $\square$

The following corollary is easily gleaned from the proof of Theorem 1.

Corollary 1. *Suppose that $\{P_i : i \in I\}$ is a family of cones in a Minkowski space $(\mathbb{R}^d, \|\cdot\|)$ satisfying*

$$\bigcup_{i \in I} (P_i \cup -P_i) = \mathbb{R}^d, \quad (1)$$

and

$$\forall i \in I \forall \text{ distinct } x, y \in P_i, \text{ if } \|x\| = \|y\| \text{ then } \pm(x - y) \notin P_i. \quad (2)$$

Then a k -distance set in $(\mathbb{R}^d, \|\cdot\|)$ has cardinality at most $(k+1)^{\#I}$. $\square$

Lemma 2. *Let S be a k -distance set in a metric space (X, ρ) with distances $\rho_1 < \rho_2 < \dots < \rho_k$. If $\rho_k/\rho_1 > 2^{k-1}$, then for some $i = 1, \dots, k-1$, the relation*

$$x \sim_i y \iff \rho(x, y) \leq \rho_i$$

is an equivalence relation.

Proof. The relation $\sim_i$ is reflexive and symmetric. If it is not transitive, there exist $x, y, z \in S$ such that $\rho(x, y), \rho(y, z) \leq \rho_i$ and $\rho(x, z) > \rho_i$. Thus $\rho_{i+1} \leq \rho(x, z) \leq \rho(x, y) + \rho(y, z) \leq 2\rho_i$. If this holds for all $i = 1, \dots, k-1$, we obtain $\rho_k \leq 2^{k-1}\rho_1$. $\square$ $\square$

Lemma 3. *The cardinality of a k -distance set in a d -dimensional Minkowski space is at most 2^{kd} .*

Proof. Let $\{x_1, \dots, x_m\}$ be a k -distance set with distances $\rho_1 < \rho_2 < \dots < \rho_k$. Set $V := \bigcup_{i=1}^m B(x_i, \rho_1/2)$. Then we have

$$\text{vol}(V) = m(\rho_1/2)^d \text{vol}(B). \quad (3)$$

Also, $V - V \subseteq B(0, \rho_k + \rho_1)$, since if $x, y \in V$, there exist i and j such that $\|x - x_i\| \leq \rho_1/2, \|y - x_j\| \leq \rho_1/2$. Thus

$$\|x - y\| \leq \|x - x_i\| + \|x_i - x_j\| + \|x_j - y\| \leq \rho_1 + \rho_k.$$

Therefore,

$$\text{vol}(V - V) \leq (\rho_1 + \rho_k)^d \text{vol}(B). \quad (4)$$

Substituting (3) and (4) into the Brunn-Minkowski inequality

$$\text{vol}(V - V)^{1/d} \geq \text{vol}(V)^{1/d} + \text{vol}(-V)^{1/d}, \quad (5)$$

we obtain $\rho_1 + \rho_k \geq m^{1/d}\rho_1$, and $m \leq (1 + \rho_k/\rho_1)^d$.

If $1 + \rho_k/\rho_1 \leq 2^k$, there is nothing to prove. Otherwise, $\rho_k/\rho_1 > 2^k - 1 \geq 2^{k-1}$, and by Lemma 2, $x \sim_i y \iff \rho(x, y) \leq \rho_i$ is an equivalence relation for some $i = 1, \dots, k-1$. By induction on k we obtain that each equivalence class, being




FIGURE 1.

an i -distance set, has at most 2^{id} points. By choosing a representative from each equivalence class, we obtain a $(k - i)$ -distance set with at most $2^{(k-i)d}$ points. Therefore, $m \leq 2^{id} 2^{(k-i)d} = 2^{kd}$. $\square$

In the proof of Theorem 3, we need the following geometric lemma, which is a modification of [25, corollary 3.2.6] in 2 dimensions.

Lemma 4. *Let B_1 be the convex hull of $\{(\pm 1, 0), (0, \pm 1)\}$ and B_∞ the square $[-1, 1]^2$. For any symmetric convex disc C in $\mathbb{R}^2$ there exists an invertible linear transformation taking C to C' such that $B_1 \subseteq C' \subseteq B_\infty$ and such that any straight-line segment contained in the boundary of C' lies completely in one of the four coordinate quadrants.*

Proof. We consider all triangles with vertices $0, x, y$, where x and y are on the boundary of C . By compactness there exist x_0 and y_0 such that the area of the triangle is a maximum. Then $\{x_0 + \lambda y_0 : \lambda \in \mathbb{R}\}$ is a support line of C at x_0 , since otherwise we can replace x_0 by a point on the side of the line opposite 0 to enlarge the area of the triangle. Similarly, $\{y_0 + \lambda x_0 : \lambda \in \mathbb{R}\}$ is a support line of C at y_0 . Since C is symmetric, it follows that C is contained in the parallelogram $\{\lambda x_0 + \mu y_0 : -1 \leq \lambda, \mu \leq 1\}$. See Figure 1.

If x_0 is an interior point of a straight-line segment contained in the boundary of C , we may shift x_0 to a boundary point of such a segment, without changing the area of the triangle. Thus C is still contained in a parallelogram as above. A similar remark holds for y_0 . We now apply the linear transformation sending x_0 and y_0 to the standard unit vectors e_1 and e_2 , respectively (see Figure 2). $\square$

Proof of Theorem 3. We have to find two cones P_1 and P_2 satisfying (1) and (2) of Corollary 1. By Lemma 4 we may replace the space by an isometric space $(\mathbb{R}^2, \|\cdot\|)$ such that the unit ball B of $\|\cdot\|$ lies between B_1 and B_∞ , and such that any straight-line segment contained in the boundary of the unit ball lies completely in a quadrant of the plane.

We provisionally let P_1 be the closed first quadrant, and P_2 the closed second quadrant. See Figure 2. Then (1) is satisfied. The only way that (2) could fail is if there is a straight-line segment contained in the boundary of the unit ball parallel to either the x -axis or the y -axis, lying in P_1 or P_2 . If there is a segment in the boundary of the unit ball in P_1 parallel to the x -axis, say, we remove the positive x -axis $\{(\lambda, 0) : \lambda > 0\}$ from P_1 . If in this case there were another straight-line segment in the boundary parallel to the x -axis in P_2 , then there would be a straight-line segment in the boundary lying in the first and second quadrants, giving a contradiction. Thus we do not have to remove the negative x -axis from P_2 ,




FIGURE 2.

and (1) is still satisfied. We do the same thing for segments parallel to the y -axis, and for P_2 . In the end, the modified P_1 and P_2 satisfy (1) and (2), and we deduce $\#S \leq (k+1)^2$ from Corollary 1.

If equality holds, then the mapping $x \mapsto (h_1(x), h_2(x))$ in the proof of Theorem 1 is a bijection from S to $\{0, \dots, k\}^2$. We now denote a point $x \in S$ by $p_{i,j}$, where $(i, j) = (h_1(x), h_2(x))$.

Suppose that two of the distances $\|p_{0,i} - p_{0,0}\|$ ($i = 1, \dots, k$) are equal, say $\|p_{0,i} - p_{0,0}\| = \|p_{0,j} - p_{0,0}\|$ with $0 < i < j$. Then, since $p_{0,j} >_2 p_{0,i} >_2 p_{0,0}$, we have $p_{0,i} - p_{0,0}, p_{0,j} - p_{0,0} \in P_2$, contradicting (2).

It follows that the distances $\|p_{0,i} - p_{0,0}\|$, $i = 1, \dots, k$ are distinct, and thus are exactly the k different distances in increasing order. Similarly, the distances $\|p_{0,i} - p_{0,1}\|$, $i = 2, \dots, k$ are in increasing order. If $\|p_{0,k} - p_{0,1}\| = \rho_k$, the three points $p_{0,0}, p_{0,1}, p_{0,k}$ again contradict (2). Thus these distances are $\rho_1, \dots, \rho_{k-1}$ in increasing order, etc. In the end we find that $\|p_{0,i+1} - p_{0,i}\| = \rho_1$ for all i . Thus $\rho_k \leq k\rho_1$, by the triangle inequality. Using the Brunn-Minkowski inequality as in the proof of Lemma 3, we find that equality holds in (5) and (4), implying that for $V := \bigcup_{i=1}^{\#S} B(x_i, \rho_1/2)$ we have $V - V = B(0, \rho_k + \rho_1)$, and $V - V$ and V are homothetic. Thus V is a ball that is perfectly packed by smaller balls. By a result of [16], this implies that the unit ball is a parallelogram. $\square$ $\square$

Proof of Theorem 4. Follows from the proof of Theorem 3 in the same way that Theorem 2 follows from Theorem 1. $\square$ $\square$

Proof of Theorem 5. Lemma 3 already gives part of the theorem. For the remaining part we apply Corollary 1. In order for a cone P to satisfy (2), it is sufficient that

$$\forall a, b \in P : \text{ if } \|a\| = \|b\| = 1, \text{ then } \|a - b\| < 1. \quad (6)$$

To see this, suppose that P does not satisfy the condition in (2), i.e. there exist distinct $x, y \in P$ such that $\|x\| = \|y\|$ and $y - x \in P$. Let $a := \|x\|^{-1}x$, $b := \|y\|^{-1}y$, $c := \|y - x\|^{-1}(y - x)$, and $0 < \lambda := \|x\| / (\|y - x\| + \|x\|) < 1$. Then $a = (1 - \lambda)(a - c) + \lambda b$, and

$$1 = \|a\| \leq (1 - \lambda)\|a - c\| + \lambda\|b\| = (1 - \lambda)\|a - c\| + \lambda,$$

implying $\|a - c\| \geq 1$.

In order for (1) to be satisfied too, we need a cover of the unit sphere by sets such that, if they are extended to positive cones, are convex.

We do this with the following construction: Let $C = \{c_1, c_2, \dots, c_m\}$ be a maximal set of unit vectors satisfying $\|c_i - c_j\|, \|c_i + c_j\| \geq \frac{1}{5}$ for all $1 \leq i < j \leq m$.

Then for any unit vector x there exists i such that $\|c_i - x\| < \frac{1}{5}$ or $\|c_i + x\| < \frac{1}{5}$. For $i = 1, \dots, m$, let P_i be the cone generated by

$$Q_i := \{x \in \mathbb{R}^d : \|x\| = 1, \|c_i - x\| < \frac{1}{5}\},$$

i.e. $P_i := \{\sum_j \lambda_j x_j : \lambda_j \geq 0, x_j \in Q_i\}$. Then the P_i 's satisfy (1) by the maximality of C . Each P_i satisfies (6): Let $\sum_j \lambda_j x_j \in P_i$, where $\lambda_j \geq 0, \|x_j\| = 1, \|c_i - x_j\| < \frac{1}{5}$ and $\|\sum \lambda_j x_j\| = 1$. Then

$$\begin{aligned} \|c_i - \sum_j \lambda_j x_j\| &= \|\sum_j \lambda_j (c_i - x_j) + (1 - \sum_j \lambda_j) c_i\| \\ &< \sum_j \lambda_j / 5 - 1 + \sum_j \lambda_j \quad (\text{since } \sum_j \lambda_j \geq 1) \\ &= \frac{6}{5} \sum_j \lambda_j - 1. \end{aligned}$$

Also, since

$$1 + \sum_j \lambda_j / 5 > \|\sum_j \lambda_j x_j\| + \sum_j \|\lambda_j x_j - \lambda_j c\| \geq \sum_j \lambda_j \|c\| = \sum_j \lambda_j,$$

we obtain $\sum_j \lambda_j < \frac{5}{4}$, and $\|c_i - \sum_j \lambda_j x_j\| < \frac{6}{5} \cdot \frac{5}{4} - 1 = \frac{1}{2}$.

A volume argument gives the upper bound for $\#C$: The balls

$$B(0, \frac{9}{10}), B(\pm c_i, \frac{1}{10}), i = 1, \dots, m$$

have disjoint interiors and are contained in the ball $B(0, \frac{11}{10})$. Therefore,

$$(\frac{9}{10})^d \text{vol}(B) + 2m(\frac{1}{10})^d \text{vol}(B) \leq (\frac{11}{10})^d \text{vol}(B),$$

giving $m \leq \frac{1}{2}(11^d - 9^d)$. $\square$

ACKNOWLEDGEMENT

This paper is part of the author's PhD thesis written under supervision of Prof. W. L. Fouché at the University of Pretoria. I thank the referees as well as Graham Brightwell for their suggestions on the layout of the paper.

REFERENCES

- [1] E. Bannai and E. Bannai, *An upper bound for the cardinality of an s -distance subset in real Euclidean space*, *Combinatorica* **1** (1981), 99–102.
- [2] E. Bannai, E. Bannai, and D. Stanton, *An upper bound for the cardinality of an s -distance subset in real Euclidean space II*, *Combinatorica* **3** (1983), 147–152.
- [3] J. Beck, *On the lattice property of the plane and some problems of Dirac, Motzkin and Erdős in combinatorial geometry*, *Combinatorica* **3** (1983), 281–297.
- [4] A. Blokhuis, *Few-distance sets*, CWI Tract 7, Stichting Mathematisch Centrum, Amsterdam, 1984.
- [5] ———, *A new upper bound for the cardinality of 2-distance sets in Euclidean space*, *Ann. Discrete Math.* **20** (1984), 65–66.
- [6] A. Blokhuis and H. A. Wilbrink, *Alternative proof of Sine's theorem on the size of a regular polygon in $\mathbb{R}^n$ with the ℓ_∞ -metric*, *Discrete Comput. Geom.* **7** (1992), 433–434.
- [7] P. Brass, *On equilateral simplices in normed spaces*, To be published in *Beiträge Algebra Geom.*, 1998.
- [8] Yu. D. Burago and V. A. Zalgaller, *Geometric inequalities*, Springer-Verlag, Heidelberg, 1988.
- [9] F. Chung, *The number of different distances determined by n points in the plane*, *J. Comb. Theory, Ser. A* **36** (1984), 342–354.
- [10] F. Chung, E. Szemerédi, and W. Trotter, Jr., *The number of distinct distances determined by a set of points in the Euclidean plane*, *Discrete Comput. Geom.* **7** (1992), 1–11.
- [11] K. Clarkson, H. Edelsbrunner, L. Guibas, M. Sharir, and E. Welzl, *Combinatorial complexity bounds for arrangements of curves and surfaces*, *Discrete Comp. Geom.* **5** (1990), 99–160.
- [12] H. T. Croft, K. J. Falconer, and R. K. Guy, *Unsolved Problems in Geometry*, Springer-Verlag, New York, 1991.

- [13] P. Erdős, *On sets of distances of n points*, Amer. Math. Monthly **53** (1946), 248–250.
- [14] ———, *On sets of distances of n points in Euclidean space*, Publ. Math. Inst. Hungar. Acad. Sci **5** (1960), 165–169.
- [15] M. Golomb, *Advanced problems and solutions. Isosceles n -points*, Amer. Math. Monthly **55** (1948), 513–514.
- [16] H. Groemer, *Abschätzungen für die Anzahl der konvexen Körper, die einen konvexen Körper berühren*, Monatsh. Math. **65** (1961), 74–81.
- [17] R. K. Guy and R. B. Kusner, *An olla-podrida of open problems, often oddly posed*, Amer. Math. Monthly **90** (1983), 196–199.
- [18] H. Harborth and L. Piepmeyer, *Three distinct distances in the plane*, Geom. Dedicata **61** (1996), 315–327.
- [19] L. M. Kelly, *Elementary problems and solutions. Isosceles n -points*, Amer. Math. Monthly **54** (1947), 227–229.
- [20] G. R. Lawlor and F. Morgan, *Paired calibrations applied to soap films, immiscible fluids, and surfaces and networks minimizing other norms*, Pacific J. Math. **166** (1994), 55–82.
- [21] F. Morgan, *Minimal surfaces, crystals, networks, and undergraduate research*, Math. Intelligencer **14** (1992), 37–44.
- [22] J. Pach and P. K. Agarwal, *Combinatorial Geometry*, John Wiley & Sons, New York, 1995.
- [23] C. M. Petty, *Equilateral sets in Minkowski spaces*, Proc. Amer. Math. Soc. **29** (1971), 369–374.
- [24] L. A. Székely, *Crossing numbers and hard Erdős problems in discrete geometry*, Combinatorics, Probability & Computing **6** (1997), 353–358.
- [25] A. C. Thompson, *Minkowski Geometry*, Encyclopedia of Mathematics and its Applications 63, Cambridge University Press, 1996.

DEPARTMENT OF MATHEMATICS AND APPLIED MATHEMATICS, UNIVERSITY OF PRETORIA, PRETORIA 0002, SOUTH AFRICA, KONRAD@MATH.UP.AC.ZA